arXiv:1206.2280v2 [math.NT] 3 Nov 2012

THE FROBENIUS-EULER FUNCTION AND ITS APPLICATIONS

SERKAN ARACI, **DEYAO GAO**, AND **MEHMET ACIKGOZ**

Abstract. In the present paper, we deal with Fourier-transformation of Frobenius-Euler polynomials. We shall give its applications by using infinite series. Our applications possess interesting properties which we state in this paper.

2010 Mathematics Subject Classification. 11S80, 11B68.

Keywords and phrases. Frobenius-Euler numbers and polynomials, Fourier transformation, infinite series.

1. **Introduction**

The ordinary Frobenius-Euler numbers are defined by means of the following generating function:

$$\sum_{n=0}^{\infty} H_n(u) \frac{t^n}{n!} = e^{H(u)t} = \frac{1-u}{e^t - u}. \tag{1.1}$$

where, in the umbral calculus, $H^n(u)$ is symbolically replaced by $H_n(u)$ in the formal series expansion of

$$e^{tH(u)} = \sum_{n=0}^{\infty} H_n(u) \frac{t^n}{n!}.$$

From expression of this definition, we state the following

$$(H(u)+1)^n - uH_n(u) = \begin{cases} 1-u & \text{if } n=0, \\ 0 & \text{if } n \in \mathbb{N}, \end{cases} \tag{1.2}$$

where $\mathbb{N}$ denotes the set of positive integers.

From (1.2), we note that

$$H_0(u) = 1, H_1(u) = -\frac{1}{1-u}, H_2(u) = \frac{1+u}{(1-u)^2}, \cdots .$$

The Frobenius Euler polynomials are also introduced as

$$e^{xt}\frac{1-u}{e^t-u} = \sum_{n=0}^{\infty} \frac{t^n}{n!} H_n(x,u). \tag{1.3}$$

By (1.1) and (1.3), we can find the following

$$H_n(x,u) = \sum_{l=0}^{n} \binom{n}{l} x^{n-l} H_l(u).$$

By expression of (1.2), it is not difficult to show that the recurrence relation for the Frobenius-Euler numbers as follows:

$$\sum_{l=0}^{n}\binom{n}{l}H_l\left(u\right)-uH_n\left(u\right)=\left(1-u\right)\delta_{0,n} \tag{1.4}$$

where $\delta_{m,n}$ is the Kronecker delta, is defined by

$$\delta_{m,n}=\begin{cases}1, & \text{if } m=n\\ 0, & \text{if } m\neq n.\end{cases}$$

Thus, we easily procure the following:

$$H_n\left(1,u\right)-uH_n\left(u\right)=0 \ \ (\text{for } n\in\mathbb{N}). \tag{1.5}$$

Thus, we arrive the following lemma.

Lemma 1. *For $n\in\mathbb{N}$, we have*

$$H_n\left(1,u\right)=uH_n\left(u\right).$$

Substituting $u=-1$ in the above lemma, it leads to

$$H_n\left(-1,u\right)=E_n\left(1\right)=-E_n$$

where E_n is called Euler numbers, as is well-known, Euler numbers are defined by the following generating function:

$$\sum_{n=0}^{\infty}E_n\frac{t^n}{n!}=\frac{2}{e^t+1}$$

(for more informations on this subjects, see[1-20]).

Recently, Fourier transformation of the special functions have been studied by many mathematicians (cf., [1], [2], [4], [13], [14], [16]). In [16], Luo gave Fourier expansions of Apostol-Bernoulli and Apostol-Euler polynomials and derived some integral representations of Apostol-Bernoulli and Apostol-Euler polynomials by using Fourier expansions. After, Bayad [1] introduced as theoretical identities of the Fourier transformation of the Apostol-Bernoulli, Apostol-Euler and Apostol-Genocchi polynomials. Next, T. Kim also defined the Euler function which is Fourier transformation of Euler polynomials. We easily see that Kim's method is different from Bayad and Luo. Actually, Kim's paper [3, pp. 131-136] motivated us to write this paper. Thus, we also give Fourier transformation of Frobenius-Euler function by using Kim's method. In this paper, we also show that this function is related to Lerch trancendent $\Phi\left(z,s,a\right)$.

2. On the Frobenius-Euler function

In this section, we consider Frobenius-Euler function by using infinite series. For $m\in\mathbb{N}$, the Fourier transformation of Frobenius-Euler function is introduced as

$$H_m\left(x,u\right)=\sum_{n=-\infty}^{\infty}a_n^{(m)}\left(u\right)e^{(2n+1)\pi ix}, \ \left(a_n^{(m)}\left(u\right)\in\mathbb{C}\right) \tag{2.1}$$

where $\mathbb{C}$ denotes the set of complex numbers and

$$a_n^{(m)}\left(u\right)=\int_0^1 H_m\left(x,u\right)e^{-(2n+1)ix\pi}dx. \tag{2.2}$$

By applying some technical method on (2.2), we procure the following

$$\begin{aligned} a_n^{(m)}(u) &= \left[\frac{H_{m+1}(x,u)}{m+1}e^{-(2n+1)ix\pi}\right]_0^1 - \frac{(2n+1)\pi i}{m+1}\int_0^1 H_{m+1}(x,u)\,e^{-(2n+1)ix\pi}dx \\ &= -\frac{u+1}{m+1}H_{m+1}(u) - \frac{(2n+1)\pi i}{m+1}a_n^{(m+1)}(u). \end{aligned}$$

So from above, it leads to the following

$$a_n^{(m+1)}(u) = \left[a_n^{(m)} + (u+1)\frac{H_{m+1}(u)}{m+1}\right]\frac{m+1}{((2n+1)\pi i)}.$$

By continuing this process, becomes as follows:

$$a_n^{(m)}(u) = \left[a_n^{(1)}(u) + (u+1)\frac{H_2(u)}{2}\right]\frac{m!}{((2n+1)\pi i)^{m-1}} \tag{2.3a}$$

$$+ (u+1)\left[\frac{1}{(2n+1)\pi i}H_m(u) + \frac{m}{((2n+1)\pi i)^2}H_{m-1}(u) + ... + \frac{m!}{4!\,((2n+1)\pi i)^{m-3}}\right].$$

We want to note that

$$\lim_{u\to -1} a_n^{(m)}(u) = a_n^{(m)}(-1) := a_n^{(m)}$$

where $a_n^{(m)} \in \mathbb{C}$ is defined by Kim in [3] as follows:

$$a_n^{(m)} = \frac{m!}{((2n+1)\pi i)^{m-1}}a_n^{(1)}(u).$$

By using (2.1) and (2.3a), we readily derive the following

$$H_m(x,u) = \sum_{n=-\infty}^{\infty}\left\{\begin{array}{c} \left[a_n^{(1)}(u) + (u+1)\frac{H_2(u)}{2}\right]\frac{m!}{((2n+1)\pi i)^{m-1}} + (u+1) \\ \times\left[\frac{1}{(2n+1)\pi i}H_m(u) + \frac{m}{((2n+1)\pi i)^2}H_{m-1}(u) + ... + \frac{m!}{4!((2n+1)\pi i)^{m-3}}\right] \end{array}\right\}e^{(2n+1)\pi i x}.$$

From this, we can state the following

$$H_m(x,u) = \sum_{n=-\infty}^{\infty}\left[a_n^{(1)}(u) + (u+1)\frac{H_2(u)}{2}\right]\frac{m!e^{(2n+1)\pi i x}}{((2n+1)\pi i)^{m-1}}$$

$$+ (u+1)\sum_{n=-\infty}^{\infty}\left[\frac{1}{(2n+1)\pi i}H_m(u) + \frac{m}{((2n+1)\pi i)^2}H_{m-1}(u) + ... + \frac{m!}{4!\,((2n+1)\pi i)^{m-3}}H_4(u)\right]e^{(2n+1)\pi i x}$$

After some calculations on the above equation, we have the following

$$\begin{aligned} H_m(x,u) &= \sum_{n=-\infty}^{\infty}\left[\frac{u+1}{u-1}\frac{1}{(2n+1)\pi i} + \frac{2}{((2n+1)\pi i)^2} + \frac{1}{2}\left(\frac{u+1}{1-u}\right)^2\right]\frac{m!e^{(2n+1)\pi i x}}{((2n+1)\pi i)^{m-1}} \\ &\quad + (u+1)\sum_{n=-\infty}^{\infty}\left[\sum_{k=0}^{m-4}\frac{1}{((2n+1)\pi i)^{k+1}}\frac{d^k}{dx^k}H_m(x,u)\mid_{x=0}\right]e^{(2n+1)\pi i x} \end{aligned}$$

As a result, we conclude the following theorem.

Theorem 1. *For $m \in \mathbb{N}$ and $0 \leq x < 1$, we have*

$$H_m(x,u) = m! \sum_{n=-\infty}^{\infty} \left[\frac{u+1}{u-1} \frac{1}{(2n+1)\pi i} + \frac{2}{((2n+1)\pi i)^2} + \frac{1}{2}\left(\frac{u+1}{1-u}\right)^2 \right] \frac{e^{(2n+1)\pi i x}}{((2n+1)\pi i)^{m-1}}$$
$$+ (u+1) \sum_{k=0}^{m-4} \frac{1}{(2\pi i)^{k+1}} \frac{d^k}{dx^k} H_m(x,u) \mid_{x=0} \sum_{n=-\infty}^{\infty} \frac{e^{(2n+1)\pi i x}}{\left(n+\frac{1}{2}\right)^{k+1}}.$$

Considering generating functions of Euler and Frobenius-Euler polynomials, we reach the following corollary.

Corollary 1. *Taking $u = -1$, we have Fourier transformation of Euler function, which is defined by Kim in* [3] *as follows:*

$$H_m(x,-1) = E_m(x) = 2m! \sum_{n=-\infty}^{\infty} \frac{e^{(2n+1)\pi i x}}{((2n+1)\pi i)^{m+1}} \quad (\text{for } 0 \leq x < 1).$$

The Lerch trancendent $\Phi(z,s,a)$ is the analytic continuation of the series

$$\Phi(z,s,a) = \sum_{n=0}^{\infty} \frac{z^n}{(n+a)^s} \tag{2.5}$$

which converges for $a \in \mathbb{C} \backslash \mathbb{Z}_0^-$, $s \in \mathbb{C}$ when $|z| < 1$; $\Re(s) > 1$ when $|z| = 1$ where $\mathbb{Z}_0^- = \mathbb{Z}^- \cup \{0\}$, $\mathbb{Z}^- = \{-1,-2,-3,...\}$. Lerch trancendent $\Phi(z,s,a)$ is the proportional not only Riemann zeta funtion, Hurwitz zeta function, the Dirichlet's eta function but also Dirichlet beta function, the Legendre chi function, the polylogarithm, the Lerch zeta function (for details, see [15], [17]).

We now want to indicate that $\sum_{n=-\infty}^{\infty} \frac{e^{(2n+1)\pi i x}}{((2n+1)\pi i)^m}$ is closely related to Lerch trancendent $\Phi(z,s,a)$. So, we compute as follows:

$$\begin{aligned} \sum_{n=-\infty}^{\infty} \frac{e^{(2n+1)\pi i x}}{((2n+1)\pi i)^m} &= \frac{1}{(2\pi i)^m} \sum_{n=-\infty}^{\infty} \frac{e^{(2n+1)\pi i x}}{\left(n+\frac{1}{2}\right)^m} \\ &= \frac{1}{(2\pi i)^m} \sum_{n=-\infty}^{-1} \frac{e^{(2n+1)\pi i x}}{\left(n+\frac{1}{2}\right)^m} + \frac{1}{(2\pi i)^m} \sum_{n=0}^{\infty} \frac{e^{(2n+1)\pi i x}}{\left(n+\frac{1}{2}\right)^m} \end{aligned}$$

After some applications on the above equation, we procure the following

$$\sum_{n=-\infty}^{\infty} \frac{e^{(2n+1)\pi i x}}{((2n+1)\pi i)^m} = -\frac{e^{\pi i x}}{(\pi i)^m} + \frac{(-1)^m e^{\pi i x}}{(2\pi i)^m} \Phi\left(e^{-2\pi i x}, m, -\frac{1}{2}\right). \tag{2.6}$$

By using Theorem 1 and (2.6), we give the following theorem.

Theorem 2. *The following equality holds true:*

$$\begin{aligned}H_m(x,u) &= m!\frac{u+1}{u-1}\left(-\frac{e^{\pi ix}}{(\pi i)^m}+\frac{(-1)^m e^{\pi ix}}{(2\pi i)^m}+\Phi\left(e^{-2\pi ix},m,-\frac{1}{2}\right)\right)\\ &+2m!\left(-\frac{e^{\pi ix}}{(\pi i)^{m+1}}+\frac{(-1)^{m+1} e^{\pi ix}}{(2\pi i)^{m+1}}+\Phi\left(e^{-2\pi ix},m+1,-\frac{1}{2}\right)\right)\\ &+\frac{1}{2}\left(\frac{u+1}{u-1}\right)^2\left(-\frac{e^{\pi ix}}{(\pi i)^{m-1}}+\frac{(-1)^{m-1} e^{\pi ix}}{(2\pi i)^{m-1}}+\Phi\left(e^{-2\pi ix},m-1,-\frac{1}{2}\right)\right)\\ &+(u+1)\sum_{k=0}^{m-4}\frac{d^k}{dx^k}H_m(x,u)\mid_{x=0}\left(-\frac{e^{\pi ix}}{(\pi i)^{k+1}}+\frac{(-1)^{k+1} e^{\pi ix}}{(2\pi i)^{k+1}}+\Phi\left(e^{-2\pi ix},k+1,-\frac{1}{2}\right)\right).\end{aligned}$$

For $u=-1$ on the above theorem, we have the following corollary.

Corollary 2. *The following identity*

$$E_m(x)=2m!\left(-\frac{e^{\pi ix}}{(\pi i)^{m+1}}+\frac{(-1)^{m+1} e^{\pi ix}}{(2\pi i)^{m+1}}+\Phi\left(e^{-2\pi ix},m+1,-\frac{1}{2}\right)\right)$$

is true.

Setting $x=1$ in Theorem 1, we obtain

(2.7)

$$\begin{aligned}H_m(1,u) &= -m!\sum_{n=-\infty}^{\infty}\left[\frac{u+1}{u-1}\frac{1}{(2n+1)\pi i}+\frac{2}{((2n+1)\pi i)^2}+\frac{1}{2}\left(\frac{u+1}{1-u}\right)^2\right]\frac{1}{((2n+1)\pi i)^{m-1}}\\ &-(u+1)\sum_{k=0}^{m-4}\frac{1}{(2\pi i)^{k+1}}\frac{d^k}{dx^k}H_m(x,u)\mid_{x=0}\sum_{n=-\infty}^{\infty}\frac{1}{\left(n+\frac{1}{2}\right)^{k+1}}.\end{aligned}$$

By expressions of (2.7) and Lemma 1, we easily see the following corollary.

Corollary 3. *The following identity holds true:*

$$\begin{aligned}uH_m(u) &= -m!\sum_{n=-\infty}^{\infty}\left[\frac{u+1}{u-1}\frac{1}{(2n+1)\pi i}+\frac{2}{((2n+1)\pi i)^2}+\frac{1}{2}\left(\frac{u+1}{1-u}\right)^2\right]\frac{1}{((2n+1)\pi i)^{m-1}}\\ &-(u+1)\sum_{k=0}^{m-4}\frac{1}{(2\pi i)^{k+1}}\frac{d^k}{dx^k}H_m(x,u)\mid_{x=0}\sum_{n=-\infty}^{\infty}\frac{1}{\left(n+\frac{1}{2}\right)^{k+1}}.\end{aligned}$$

Now, by using Kim's method in [3], we discover the following

$$\begin{aligned}\text{(2.8a)}\qquad \frac{1}{1-ue^{-t}} &= \sum_{n=0}^{\infty}u^n e^{-nt}=\sum_{n=0}^{\infty}u^n\left(e^{-t}\right)^n=\sum_{n=0}^{\infty}u^n\left(\sum_{k=0}^{\infty}(-1)^k\frac{t^k}{k!}\right)^n\\ &=\sum_{n=0}^{\infty}\left(\sum_{a_1+a_2+\ldots+a_n=n}\frac{n!}{(a_1)!\,(a_2)!\ldots}\frac{(-1)^{a_1+2a_2+\ldots}}{(1!)^{a_1}(2!)^{a_2}\ldots}\right)t^{a_1+2a_2+\ldots}\end{aligned}$$

Let $p(i,j) : a_1 + 2a_2 + ... = i, a_1 + a_2 + ... = j$, from expression of (2.8a), we compute as follows:

$$\frac{1}{1-ue^{-t}} = \sum_{m=0}^{\infty}\left(\sum_{n=0}^{m} u^n \sum_{p(m,n)} \frac{n!}{(a_1)!\,(a_2)!...\,(a_m)!}\frac{(-1)^{a_1+2a_2+...+ma_m}}{(1!)^{a_1}\,(2!)^{a_2}\,...\,(m!)^{a_m}}\right) t^{a_1+2a_2+...+ma_m}$$

$$= \sum_{m=0}^{\infty}(-1)^m\left(\sum_{n=0}^{m} n!u^n \sum_{p(m,n)} \frac{m!}{(a_1)!\,(a_2)!...\,(a_m)!}\frac{(-1)^{m}}{(1!)^{a_1}\,(2!)^{a_2}\,...\,(m!)^{a_m}}\right)\frac{t^m}{m!}$$

$$= \sum_{m=0}^{\infty}\left[(-1)^m\sum_{n=0}^{m} n!u^n s_2(m,n)\right]\frac{t^m}{m!}. \tag{2.9a}$$

where $s_2(m,n)$ is the second kind stirling number.

Via the definition of Frobenius-Euler numbers, we readily derive the following

$$\begin{aligned}\frac{1}{1-ue^{-t}} &= \frac{u^{-1}}{u^{-1}-1}\frac{1-u^{-1}}{e^{-t}-u^{-1}} \\ &= \frac{1}{1-u}\sum_{m=0}^{\infty}(-1)^m H_m\left(u^{-1}\right)\frac{t^m}{m!}.\end{aligned} \tag{2.10}$$

By comparing the coefficients of $\frac{t^n}{n!}$ on the both sides of (2.9a) and (2.10), we reach the following theorem.

Theorem 3. *The following equality holds true:*

$$\frac{1}{1-u}H_m\left(u^{-1}\right) = \sum_{n=0}^{m} n!u^n s_2(m,n).$$

Corollary 4. *Substituting $u = -1$ in the above theorem, we get the following, which is defined by Kim* [3]

$$E_m = 2\sum_{n=0}^{m} n!\,(-1)^n s_2(m,n).$$

REFERENCES

[1] A. Bayad, Fourier expansions for Apostol-Bernoulli, Apostol-Euler and Apostol Genocchi polynomials, Mathematics of Computation, Volume 80, Number 276, October 2011, Pages 2219–2221.

[2] J. Choi, D. S. Kim, T. Kim and Y. H. Kim, A note on Some identities of Frobeniu-Euler Numbers and Polynomials, International Journal of Mathematics and Mathematical Sciences, Volume 2012, Article ID 861797, 9 pages.

[3] T. Kim, Note on the Euler numbers and polynomials, Advanced Studies in Contemporary Mathematics, vol. 17, no. 2, pp. 131–136, 2008.

[4] T. Kim and B. Lee, Some Identities of the Frobenius-Euler polynomials, Abstract and Applied Analysis, Volume 2009, Article ID 639439, 7 pages.

[5] T. Kim, Euler Numbers and Polynomials Associated with Zeta Functions, Abstract and Applied Analysis 2008 (2008), Article ID 581582, 11 pages.

[6] T. Kim, q-Volkenborn integration, Russ. J. Math. Phys. 9 (2002), 288-299.

[7] T. Kim, q-Bernoulli numbers and polynomials associated with Gaussian binomial coefficients, Russ. J. Math. Phys. 15 (2008) , 51-57.

[8] T. Kim, q-extension of the Euler formula and trigonometric functions, Russian J. Math. Phys. 14 (2007), 275-278.

[9] T. Kim, On Euler-Barnes multiple zeta functions, Russian J. Math. Phys. 10 (2003), 261-267.

[10] T. Kim, q-Euler numbers and polynomials associated with p-adic q-integrals, J. Nonlinear Math. Phys., 14 (2007), no. 1, 15–27.
[11] T. Kim, New approach to q-Euler polynomials of higher order, Russ. J. Math. Phys., 17 (2010), no. 2, 218–225.
[12] T. Kim, Some identities on the q-Euler polynomials of higher order and q-Stirling numbers by the fermionic p-adic integral on $\mathbb{Z}_p$, Russ. J. Math. Phys., 16 (2009), no.4, 484–491.
[13] T. Kim, Identities involving Frobenius-Euler polynomials arising from non-linear differential equatiaons, arXiv:1201.5088v1 [math.NT].
[14] Y. Simsek, Y. Osman, V. Kurt, On interpolation functions of the twisted generalized Frobenius-Euler numbers, Adv. Stud. Contemp. Math. 15 (2007), 187-194.
[15] H. M. Srivastava and J. Choi, Series Associated with the Zeta and Related Functions, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2001.
[16] Q-M. Luo, Fourier expansions and integral representations for the Apostol-Bernoulli and Apostol-Euler polynomials, Math. Comp., Volume 78 (2009), No. 268, 2193–2208.
[17] M. Acikgoz and Y. Simsek, On multiple interpolation functions of the Nörlund-Type q-Euler polynomials, Abstract and Applied Analysis, Vol. 2009, Article ID 382574, 14 pages.
[18] S. Araci, M. Acikgoz, K. H. Park and H. Jolany, On the unification of two families of multiple twisted type polynomials by using p-adic q-integral on $\mathbb{Z}_p$ at $q = -1$, to appear in Bulletin of the Malaysian Mathematical Sciences and Society.
[19] S. Araci, D. Erdal and J. J. Seo, A study on the fermionic p-adic q-integral representation on $\mathbb{Z}_p$ associated with weighted q-Bernstein and q-Genocchi polynomials, Abstract and Applied Analysis, Volume 2011, Article ID 649248, 10 pages.
[20] S. Araci, M. Acikgoz and K. H. Park, A note on the q-analogue of Kim's p-adic log gamma type functions associated with q-extension of Genocchi and Euler numbers with weight α, Bulletin of the Korean Mathematical Society (accepted for publication).

University of Gaziantep, Faculty of Science and Arts, Department of Mathematics, 27310 Gaziantep, TURKEY
E-mail address: `mtsrkn@hotmail.com`

Yuren Lab., No. 8 Tongsheng Road, Changsha, P. R. China
E-mail address: `13607433711@163.com`

University of Gaziantep, Faculty of Science and Arts, Department of Mathematics, 27310 Gaziantep, TURKEY
E-mail address: `acikgoz@gantep.edu.tr`